



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/66, 9/20, 9/48	A1	(11) International Publication Number: WO 00/21540 (43) International Publication Date: 20 April 2000 (20.04.00)
(21) International Application Number: PCT/EP99/07287 (22) International Filing Date: 1 October 1999 (01.10.99) (30) Priority Data: 98119102.6 9 October 1998 (09.10.98) EP (71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH]; Grenzacherstrasse 124, CH-4070 Basle (CH). (72) Inventors: GABEL, Rolf-Dieter; Kurpfalzring 96, D-68723 Schwetzingen (DE). PREIS, Walter; Mandelring 70, D-67433 Neustadt (DE). WOOG, Heinrich; Lindenstrasse 6, D-69514 Laudenbach (DE). (74) Agent: WITTE, Hubert; Grenzacherstrasse 124, CH-4070 Basle (CH).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: COMPOSITIONS CONTAINING DIPHOSPHONIC ACIDS (57) Abstract <p>The invention relates to a solid pharmaceutical form of administration containing a diphosphonic acid or a physiologically compatible salt thereof as the active substance, wherein the active substance is present in granulate form, optionally together with pharmaceutical adjuvants, in the inner phase and the outer phase contains a lubricant in the form of less than 5 % by weight of stearic acid relative to the total weight of the form of administration.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Compositions containing diphosphonic acids

The invention relates to a solid pharmaceutical form of
5 administration containing a diphosphonic acid or a
physiologically compatible salt thereof as the active
substance and stearic acid as lubricant in the outer
phase. The invention also relates to a process for
preparing the said form of administration.

10

Pharmaceutical forms of administration of diphosphonic
acids are known for treatment of calcium metabolism
diseases. Drugs containing these active substances are
used for treating hypercalcaemia and also for treating
15 tumour osteolysis resulting from bone metastases. They
can also be successfully used for treatment of
osteoporosis and resulting pain.

Since the active substances for treating diseases of this
20 kind frequently have to be administered over a long
period, oral application is very advantageous since it is
usually more acceptable by the patient.

Oral forms of administration are known in the case of
25 some diphosphonic acids and salts thereof. For example
EP-B 0 275 468, EP-B 0 625 355 (both Boehringer Mannheim)
and WO 93/21907 (Leiras Oy) disclose pharmaceutical
preparations containing clodronic acid (dichloromethylene
diphosphonic acid) or salts thereof. WO 93/09785
30 (Procter & Gamble Pharmaceuticals) discloses oral forms
of administration of risedronate (the salt of 3-pyridyl-
1-hydroxyethylidene-1,1-diphosphonic acid). WO 93/21907
and WO 93/09785 describe oral forms of administration
provided with a coating which dissolves only at pH above

5 or 5.5. The aim is to ensure that the forms of administration travel through the stomach and the active principle is released only in the intestinal tract.

- 5 The solid forms of administration of diphosphonic acids or salts thereof described in the prior art contain the active substance and selected pharmaceutical adjuvants, with which the active principle must be compatible, in the inner phase and selected pharmaceutical adjuvants in
10 the outer phase, more particularly for ensuring that the preparation can be easily processed in a capsule-filling machine or tablet press. For example EP-B 0 275 468 describes clodronate-containing drugs with a high proportion of 80 - 95% active substance, a filler content
15 of 2 - 10% and a lubricant content of 0 - 5% in the granulate, to which is added an outer phase in the form of a lubricant, preferably magnesium stearate and talcum, in a proportion of 1 - 5%.
- 20 During the development of a capsule or tablet or other solid form of administration, special attention is usually paid to the adjuvants in the outer phase.

The selection and proportion of a suitable lubricant in
25 the outer phase is particularly important, since it has great influence on the physical properties of the forms of administration under development. The choice and proportion determine whether the substance filling the capsule or tablet can be processed without difficulty on
30 a suitable machine over a prolonged period or whether the tablets stick to the compression moulding dies in the machine. Sufficient lubricant must therefore be added to the outer phase. If however the proportion of lubricant is too high, there may be other adverse effects. For

example the granulate may become so water-repellent that the resulting drug disintegrates only slowly and the desired dissolution rate (practically complete release of the active substance after 30 minutes) is not reached.

5

The following known lubricants can be used in the outer phase: magnesium stearate, calcium stearate, talcum, sodium stearyl fumarate, macrogol or hydrogenated esters of fatty acids with glycerine and stearic acid.

10

For example in EP-B 0 275 468, which describes oral forms of administration for clodronate, magnesium stearate and talcum are together used as a lubricant in the outer phase. EP-B 0 625 355 (Boehringer Mannheim GmbH)

15 discloses magnesium stearate as the only lubricant in the outer phase of clodronate forms of administration. WO 93/21907 (Leiras Oy, clodronate) Example 1, describes the use of talcum and magnesium stearate as a lubricant in the outer phase and stearic acid as a lubricant in the
20 inner phase. WO 93/09785 (Procter & Gamble, risedronate) Example 3, discloses stearic acid lubricant in a proportion of 5.8% by weight relative to the tablet core.

It has been found, however, that particularly when the
25 proportions of active substance are low, the lubricant or the concentrations thereof are not optimum, since dissolution rates of 85% after 30 minutes, indicating uniform and almost complete release of the active substance, are not obtained or else the dissolution rates
30 fall rapidly after stress through heating above room temperature.

The object of the invention therefore is to develop a pharmaceutical form of administration in which the active

substances are diphosphonic acids or physiologically compatible salts thereof and which is stable enough for the active substance to be released uniformly and almost completely within 30 minutes and for no reduction in the rate of dissolution to occur even after temperature stress. This should apply both to high and low contents of active substance in the form of administration.

Surprisingly it has been found that solid forms of administration containing less than 5% by weight of stearic acid lubricant in the outer phase, relative to the total weight of the form of administration, e.g. 0.1 to 4.9% by weight, have dissolution rates of at least 85% after 30 minutes, and the rates do not change even after weeks of exposure to temperatures of 40 - 50°C. This applies both to low and to high contents of active substance in the form of administration.

Less than 5% by weight stearic acid in the outer phase, relative to the total weight of the form of administration, results on the one hand in a sufficient lubricant effect, so that the tablet or capsule filler does not stick to the processing machines, and on the other hand the granulated active substance does not become water-repellent.

This aspect of the invention therefore relates to solid pharmaceutical forms of administration in which the active substance is a diphosphonic acid or a physiologically compatible salt thereof, wherein the active substance in granulate form, optionally together with pharmaceutical adjuvants, is present in the inner phase and the outer phase contains a lubricant in the form of stearic acid in proportions of less than 5% by

weight relative to the total weight of the form of administration.

The outer phase preferably contains stearic acid in a proportion of 0.1 to 3%, particularly 0.9 to 3% by weight, relative to the total weight of the form of administration. Particularly preferably stearic acid is added in a proportion of 1.5 to 2.7% by weight relative to the total weight of the form of administration, in which case the rate of release will be at least 90% (determined by the Paddle method after the USP).

The granulated active substance can contain pharmaceutically acceptable adjuvants and/or additives such as lactose, starch, glucose, mannitol, calcium carbonate, calcium phosphate, microcrystalline cellulose, hydroxypropyl methyl cellulose or other substances known for this purpose in the art.

The form of administration according to the invention can also contain other pharmaceutical adjuvants in the outer phase, more particularly a disintegrating agent, all known disintegrating agents being usable. More particularly cross-linked polyvinyl pyrrolidone (Crospovidone USPNF) has given good service as a disintegrating agent for the purposes of the invention.

The following bisphosphonates are active substances which can be used according to the invention in the form of free acids or pharmaceutically compatible salts or hydrates, particularly sodium salts:

(4-amino-1-hydroxybutylidene)bis-phosphonate (alendronate),

- (Dichloromethylene)bis-phosphonate (clodronate),
[1-hydroxy-3-(1-pyrrolidinyl)-propylidene]bis-phosphonate
(EB-1053),
(1-hydroxyethylidene)bis-phosphonate (etidronate),
5 [1-hydroxy-3-(methyl pentyl amino)propylidene]bis-
phosphonate (ibandronate),
[Cycloheptylamino)-methylene]bis-phosphonate
(incadronate),
(6-amino-1-hydroxyhexylidene)bis-phosphonate
10 (neridronate),
[3-(dimethylamino)-1-hydroxypropylidene]bis-phosphonate
(olpadronate),
(3-amino-1-hydroxypropylidene)bis-phosphonate
(pamidronate),
15 [1-hydroxy-2-(3-pyridinyl)ethylene]bis-phosphonate
(risedronate),
[[(4-chlorophenyl)thiol]-methylene]bis-phosphonate
(tiludronate),
[1-hydroxy-2-imidazo-(1,2-a)pyridin-3-yl ethylidene]bis-
20 phosphonate (YH 529),
[1-hydroxy-2-(1H-imidazol-1-yl)ethylidene]bis-phosphonate
(zoledronate).

- Ibandronate, etidronate, clodronate, risedronate,
25 pamidronate or alendronate or free acids thereof are
preferred active substances according to the invention.
These substances and production thereof are known and
described e.g. in the following references: US Patent No.
4 705 651 (Alendronate), US Patent No. 4 927 814
30 (Ibandronate), US Patents Nos. 3 468 935, 3 400 147,
3 475 486 (Etidronate), O.T. Quimby et al., J. Org. Chem.
32, 4111 (1967) (Clodronate), US Patent No. 4 505 321
(Risedronate) and US Patents Nos. 4 134 969 and 3 962 432
(Pamidronate).

- The proportion of active substance in the form of administration according to the invention can be up to 95% by weight relative to the total weight of the form of administration. Active substance contents of 0.2 - 30% by weight, relative to the total weight of the form of administration, are particularly preferred. The method according to the invention can particularly preferably be used to make oral forms of administration containing 0.25 - 100 mg of active substance per unit dose, particularly up to 50 mg per unit dose. The term "unit dose" denotes the discrete form of administration, i.e. the individual tablet or capsule.
- Particularly preferred according to the invention is a form of administration in which the active substance is ibandronic acid (1-hydroxy-3-(N-methyl-N-pentyl)amino-propyl-1,1-diphosphonic acid) or physiologically compatible salts thereof, e.g. the sodium salt.
- In order to prepare the form of administration according to the invention, the components are mixed dry. The active substance, preferably together with a conventional binder such as starch paste or polyvinyl pyrrolidone K25, and optionally with addition of pharmaceutically acceptable additives and adjuvants (excipients of the inner phase), is granulated wet. The wet granulate is then dried and screened.
- The outer phase is then added to the mixture. Either the components of the outer phase (stearic acid and adjuvants), are first mixed together and added to the granulate in a further step, or the stearic acid and any

other adjuvants in the outer phase are added individually and directly to the granulate.

The mixture according to the invention can easily be
5 processed using automatic equipment and then compressed to form tablets or filled into conventional gelatine capsules. Tablets so prepared may be coated with conventional films such as described, e.g., in WO 97/39755.

10

Accordingly the invention also relates to a process for producing a solid pharmaceutical form of administration in which the active substance is a diphosphonic acid or a physiologically compatible salt thereof, wherein the
15 active substance is processed by known methods with pharmaceutical adjuvants to obtain a granulate, less than 5% by weight of stearic acid lubricant is added to the resulting inner phase, and optionally further adjuvants are added to the mixture and the mixture is filled into
20 capsules or compressed to form tablets.

The tablets and capsule sizes are preferably so chosen as to give a concentration of active substance of 0.25 - 100 mg per unit dose. This determines the size of the form
25 of administration according to the invention, depending on the biological potency of the active substances and any adjuvants capable of increasing it.

The forms of administration produced according to the
30 invention, containing less than 5% by weight stearic acid in the outer phase, result in free-flowing, pourable compositions and do not adhere to the moulds or tools when compressed or filled into capsules.

In comparative tests using magnesium stearate lubricant in identical quantities in the outer phase, an in vitro release rate of 56% was found after 30 minutes. If these capsules were additionally heat-stressed at 40 - 50°C in
5 a drying cupboard for several weeks and the rate of release was measured again, the 30-minute value fell to below 30%.

The invention will be additionally described in the
10 following examples without being limited thereto.

Comparative Example 1:

Production of 5.0 mg capsules containing 1.8% by weight
15 magnesium stearate lubricant.

Item	Composition	(mg/capsule)
1	Na-Ibandronate, monohydrate	5.63
2	Lactose 200 (D80)	19.37
3	Lactose D30	249.00
4	Polyvinyl pyrrolidone K25	9.00
5	Lactose D30	128.00
6	Polyvinyl pyrrolidone, cross-linked	25.00
7	Magnesium stearate	8.00
	Weight	<u>444.00</u>

The amount of active substance is equivalent to 5.0 mg free acid.

Processing:

A preliminary mixture was made from the active substance (Item 1) and lactose 200 (Item 2). The preliminary mixture was then wet granulated with additional adjuvants
5 such as lactose D30 (Item 3), using polyvinyl pyrrolidone binder (Item 4). Additional lactose (Item 5) was then mixed with the granulate after drying and screening. The additives for the outer phase (Items 6 and 7) were then added individually to the mixture.

10

The resulting substance was filled into capsules in suitable machines. The capsules were tested as part of in-process control and immediately after production had an in vitro release rate of 56% after 30 minutes. The
15 release rate was determined by the Paddle method after the USP.

Comparative Example 2:

Production of 5.0 mg capsules containing 0.91% by weight magnesium stearate lubricant.

5

Item	Composition	(mg/capsule)
1	Na-Ibandronate, monohydrate	5.63
2	Lactose D80	19.37
3	Lactose D30	249.00
4	Polyvinyl pyrrolidone K25	9.00
5	Lactose D30	128.00
6	Polyvinyl pyrrolidone, cross-linked	25.00
7	Magnesium stearate	4.00
	Weight	<u>440.00</u>

The amount of active substance is equivalent to 5.0 mg free acid.

10 The capsules were produced as in Comparative Example 1.

The result for in vitro release after 30 minutes was 56%.

15 The capsules in Comparative Examples 1 and 2 were heat-stressed at 50°C in a drying cupboard for a number of weeks, after which the release rate was determined again. This fell to a 30-minute value below 30%.

Example 1:

Production of 5.0 mg capsules according to the invention containing 0.9 and 1.8% by weight stearic acid lubricant.

5

Item	Composition	a) (mg/capsule)	b) (mg/capsule)
1	Na-Ibandronate, monohydrate	5.63	5.63
2	Lactose D80	19.37	19.37
3	Lactose D30	249.00	249.00
4	Polyvinyl pyrrolidone K25	9.00	9.00
5	Lactose D30	128.00	128.00
6	Polyvinyl pyrrolidone, cross-linked	25.00	25.00
7	Stearic acid	(0.9%) 4.00	(1.8%) 8.00
	Weight	<u>440.00</u>	<u>444.00</u>

The capsule-filling material was produced as in Comparative Examples 1 and 2. As in these Examples, the additives 6 and 7 constituted the outer phase.

10

After drying and screening, the material was filled into size 0 capsules.

The result for in vitro release after 30 minutes was

15

- a) 90% for the batch containing 4.0 mg stearic acid and
- b) 101% for the batch containing 8.0 mg stearic acid.

The capsules in the present Example 1 were also heat-stressed at 50°C in a drying oven for a number of weeks. The rates of dissolution were then determined and were the same as before heat-stressing.

5

Example 2:

Production of 20 mg tablets according to the invention containing 2.5% by weight stearic acid lubricant.

10

Item	Composition	(mg/capsule)
1	Na-Ibandronate	21.38
2	Lactose D30	45.52
3	Hydroxypropyl methyl cellulose	2.00
4	Cellulose, microcrystalline	3.00
5	Polyvinyl pyrrolidone, cross-linked	5.50
6	Stearic acid	(2.5%) 2.00
	Weight	<u>79.40</u>

The amount of active substance is equivalent to 20.0 mg of free acid.

15 Processing:

The active substance was mixed with the adjuvants (Items 2, 3 and 4) and wet granulated with water. A mixture constituting the outer phase (Items 5 and 6) was added to
20 the granulate after drying and screening. The material

ready for compressing was then compressed to form tablets.

The resulting tablets were tested for the in vitro
5 release rate immediately after production. The value after 30 minutes was 102%.

Example 3:

- 10 Production of 50 mg tablets according to the invention containing 2.5% by weight stearic acid lubricant.

Item	Composition	(mg/capsule)
1	Na-Ibandronate	53.45
2	Lactose D30	113.80
3	Hydroxypropyl methyl cellulose	5.00
4	Cellulose, microcrystalline	7.50
5	Polyvinyl pyrrolidone, cross-linked	13.75
6	Stearic acid	(2.5%) 5.00
	Weight	<u>198.50</u>

The amount of active substance is equivalent to 50.0 mg
15 of free acid.

Processing:

The active substance was mixed with the adjuvants (Items
20 2, 3 and 4) and wet granulated with water. The constituents of the outer phase (Items 5 and 6) were

individually mixed with the granulate after drying and screening. The material ready for pressing was then compressed to form tablets.

- 5 During tests on stability at various temperatures up to 40°C, the rate of release was repeatedly determined after various time intervals. Even after 26 weeks at temperatures of 40°C, there were no observable differences from the original rate of release.

C L A I M S

1. A solid pharmaceutical form of administration
containing a diphosphonic acid or a physiologically
5 compatible salt or hydrate thereof as the active
substance, wherein the active substance is in the form of
a granulate, optionally together with pharmaceutical
adjuvants, in the inner phase and the outer phase
contains a lubricant in the form of less than 5% by
10 weight of stearic acid relative to the total weight of
the form of administration.
2. A form of administration according to claim 1 which,
in the outer phase, contains 0.1 to 3% by weight of
15 stearic acid relative to the total weight of the form of
administration.
3. A form of administration according to claim 1 which,
in the outer phase, contains 0.9 to 3% by weight of
20 stearic acid relative to the total weight of the form of
administration.
4. A form of administration according to any of claims
1 to 3, which, in the outer phase, contains 1.5 to 2.7%
25 by weight of stearic acid, relative to the total weight
of the form of administration.
5. A form of administration according to any of claims
1 to 4, wherein the active substance is ibandronate,
30 etidronate, clodronate, risedronate, pamidronate or
alendronate, or the corresponding free acid.

6. A form of administration according to any of claims 1 to 5, wherein the active substance is present in a proportion of 0.25 - 100 mg per unit dose.
- 5 7. A form of administration according to any of claims 1 to 5, wherein the active substance is present in a proportion of 0.5 - 50 mg per unit dose.
8. A form of administration according to any of claims
10 1 to 7, wherein an additional pharmaceutical adjuvant in the form of a disintegrating agent is contained in the outer phase.
9. A form of administration according to claim 8,
15 wherein the disintegrating agent is polyvinyl pyrrolidone, cross-linked (Crospovidone USPNF).
10. A process for producing a solid pharmaceutical form of administration according to claims 1 to 9, wherein the
20 active substance together with pharmaceutical adjuvants is converted by known methods into a granulate, less than 5% by weight of stearic acid lubricant and optional other adjuvants are added to the resulting inner phase, and the mixture is filled into capsules or compressed to form
25 tablets.
11. A process according to claim 10, wherein the stearic acid and any other adjutants in the outer phase are individually mixed with the granulate.
30
12. A process according to claim 10, wherein the stearic acid is first mixed with other adjuvants for the outer phase and the resulting mixture is then mixed with the granulate.

13. Pharmaceutical forms of administration and production thereof as described hereinbefore, particularly in the examples.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/07287

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/66 A61K9/20 A61K9/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 275 468 A (BOEHRINGER MANNHEIM GMBH) 27 July 1988 (1988-07-27) cited in the application page 2, line 15-21 page 2, column 40-44 page 2, line 51,52 page 3, line 5-9 examples claims	1-12
A	WO 93 21907 A (LEIRAS OY ;POSTI JUHANI (FI); KATILA KIRSI (FI); RANTALA PERTTI (F) 11 November 1993 (1993-11-11) cited in the application page 1, line 3-7 page 4, line 1-9 page 5, line 1-28 page 6, line 21-25	1-12

-/-



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

19 January 2000

Date of mailing of the international search report

26/01/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Epskamp, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/07287

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 96 41618 A (SANOFI WINTHROP INC) 27 December 1996 (1996-12-27) page 13, line 8-11 examples 3,4,6 -----</p>	1-12

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 99/07287

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0275468 A	27-07-1988	AT 60711 T	15-02-1991
		CA 1298196 A	31-03-1992
		DK 658387 A	21-06-1988
		GR 3001615 T	23-11-1992
		HK 59894 A	08-07-1994
		SG 21094 G	10-06-1994
		US 4859472 A	22-08-1989
WO 9321907 A	11-11-1993	SE 501389 C	30-01-1995
		AT 143802 T	15-10-1996
		AU 3955093 A	29-11-1993
		CA 2134196 A	11-11-1993
		CZ 9402614 A	15-02-1995
		DE 69305335 D	14-11-1996
		DE 69305335 T	30-04-1997
		DE 637236 T	06-07-1995
		DK 637236 T	17-03-1997
		EP 0637236 A	08-02-1995
		ES 2094539 T	16-01-1997
		FI 944961 A	21-10-1994
		GR 3022249 T	30-04-1997
		HK 13097 A	14-02-1997
		HU 74572 A	28-01-1997
		IL 105403 A	10-03-1998
		JP 7505898 T	29-06-1995
		NO 944001 A	21-10-1994
		SE 9201299 A	25-10-1993
		US 5525354 A	11-06-1996
WO 9641618 A	27-12-1996	US 5656288 A	12-08-1997
		AU 5980896 A	09-01-1997
		BR 9609477 A	25-05-1999
		CA 2223568 A	27-12-1996
		EP 0831789 A	01-04-1998
		JP 11507924 T	13-07-1999
		NO 975634 A	29-01-1998